WATER SUPPLY OUTLOOK





California Nevada River Forecast Center NOAA - National Weather Service Sacramento, California

DEFINITIONS:

Acre-Feet: The volume equal to one acre covered one foot deep (43,560 cubic feet).

Forecast Period: Generally, April 1st through July 31st, unless otherwise noted.

April-High Forecast Period: For the Lake Tahoe Stage Rise, the period from April 1st to the highest recorded lake stage level.

April 1st Average: The April 1st snowpack average is used as a reference point because it is normally the end of the winter snowfall season and the beginning of the spring runoff season.

Residual Period: The forecast period from the first of the current month through September 30th.

Probability Forecasts: Precipitation and snowfall accumulation of known probability as determined by analysis of past records are utilized in the preparation of probability runoff forecasts. The forecasts include an evaluation of the standard error of the prediction model. The forecasts are presented at three levels of probability as follows:

- **Most Probable Volume:** Given the current hydrometeorological conditions to date, this is the best estimate of what the actual runoff volume will be this season.
- Most Probable Volume (% Normal): Most probable volume in percent of the 1961-1990 average.
- **Reasonable Maximum Volume:** Given current hydrometeorological conditions, the seasonal runoff that has a 10 percent chance of being exceeded.
- Reasonable Minimum Volume: Given current hydrometeorological conditions, the seasonal runoff that has a 90 percent chance of being exceeded.

SNOTEL: Acronym for SNOw TELemetry. This is a automated snow measurement system operated by the USDA - Natural Resources Conservation Service. These sites use meteor burst communications technology to transmit hydrometeorological information such as snow water equivalent from snow pillows, accumulated precipitation and maximum, minimum and average air temperature.

Water equivalent: The depth of water that would result from melting the snowpack at a point.

Water Year: The period from October 1st through September 30th.

Snow basins in California got off to a good start this water year with many watersheds in the central and southern Sierra Nevada receiving average to much above average precipitation during October. However, dry weather dominated during November and December bringing seasonal averages to much below average by the end of December. Although water managers are contending with low snow amounts at the present time, conditions are partly mitigated by much above average carryover storage from last year's bountiful storms. Even though one of the wettest months in the wet season has come and gone, there is room for cautious optimism that plenty of time remains for conditions to improve.

Both December and seasonal (October 1, 2011 to December 31, 2011) precipitation was much below average for the region. It was among one of the driest Decembers for many locations in the Upper Sacramento and Sierra Nevada. To cite a few examples, it was the 3rd driest December in 123 years of record at Mount Shasta City, California. It was the driest December on record at South Lake Tahoe and Mammoth Lakes Ranger Station.

<u>Basin</u>	Dec % of Avg	WY % of Avg
Trinity	11	54
Upper Sacramento	5	39
Central Sierra	1	39
Southern Sierra	1	57
Walker	1	30
Carson	2	29
Truckee	3	33
Klamath	30	50

Electronic snow sensor readings as of January 1st, 2012 show much more meager snow pack conditions than at this time last year.


	% of Average	% of Average
<u>Basin</u>	<u>January 1, 2012</u>	January 1, 2011
Trinity/Upr Sac/Nrn Sierra	24	179
Central Sierra	14	198
Southern Sierra	27	271
Tahoe-Truckee	13	212
Carson-Walker	10	216
Humboldt	23	193
Upper Klamath	33	193


The unusually dry December resulted in much below average runoff during the month:

<u>Basin</u>	Dec % of Avg	WY % of Avg
Trinity-Sacramento	25	45
San Joaquin	17	51
Tulare Lake	55	115
East Side Sierra	49	63
Humboldt	72	86
Upper Klamath	52	65

Reservoir storage in the Upper Sacramento/Sierra Nevada reflects the wet conditions during the past year with above average reservoir contents. The key reservoirs at Shasta Lake and Lake Oroville are at 111 percent of average as of December 31. Stored water in the Sacramento region as of December 31st was at 110 percent of average for the date, the San Joaquin at 132, and the Tulare Lake watershed at about 123 percent. East-side Sierra reservoirs were at 140 percent of average. The lake level at Lake Tahoe stood at 6226.76 feet (or 3.76 feet above its natural rim altitude of 6223.0 feet) as of December 31st. Usable storage was 457,500 acre-feet or 131 percent of average. It was 187,900 acre-feet (53 percent of average) at this time last year. Storage at Lahontan Reservoir in Nevada stands at 126 percent of average as of December 31st while Rye Patch Reservoir is at 138 percent. Storage at Upper Klamath Lake is about 85 percent of average.

April through July runoff forecasts vary from 50 to 74 percent for the Upper Sacramento and west slope Sierra Nevada basins. They range from 31 to 54 percent of average (1971-2000) for the east side Sierra Nevada basins and 21 to 36 percent for forecast points on the main stem Humboldt River. The April through September forecast for the Upper Klamath Lake inflow is 58 percent.





Upper Klamath and Trinity River Basins

COASTAL BASINS

COASTAL BASINS		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
Williamson River Sprague, blo	Apr-Sep	235	61	350	175	385
Sprague River Chiloquin, nr	Apr-Sep	115	50	235	75	230
Upper Klamath Falls River Inflow	Apr-Sep	300	58	550	165	515
Lost River Gerber Reservoir Inflow Clear Lake Reservoir Inflow	Feb-Jul Feb-Jul	24 57	51 54	50 100	10.0 25	47 105
Scott River Fort Jones, nr	Apr-Jul	100	55	235	30	181
Trinity R River Trinity Lake	Apr-Jul	375	59	715	200	635
SACRAMENTO RIVER BASIN		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
SACRAMENTO RIVER ABOVE BEND BRID	OGE					
Pit River Montgomery Ck, nr	Apr-Jul	645	69	1020	510	940
Mccloud River Shasta Lk, abv	Apr-Jul	275	74	440	190	370
Sacramento River Delta Shasta Dam Bend Bridge, abv, Red Bluff, nr	Apr-Jul Apr-Jul Apr-Jul	195 1240 1600	67 69 66	385 2150 3130	70 850 1050	290 1790 2440

		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
FEATHER RIVER ABOVE OROVILLE RES	SERVOIR					
North Yuba River Goodyears Bar, blo	Apr-Jul	160	59	350	55	273*
South Yuba River Langs Crossing	Apr-Jul	140	62	290	50	225*
Yuba River Englebright Reservoir	Apr-Jul	540	54	1330	160	995
YUBA RIVER ABOVE SMARTVILLE						
North Yuba River Goodyears Bar, blo	Apr-Jul	160	59	350	55	273*
South Yuba River Langs Crossing	Apr-Jul	140	62	290	50	225*
Yuba River Englebright Reservoir	Apr-Jul	540	54	1330	160	995
AMERICAN RIVER ABOVE FOLSOM RES	ERVOIR					
Middle Fork American River Auburn, nr	Apr-Jul	265	54	590	110	490*
Silver Creek Union Valley Camino Dam, blo	Apr-Jul Apr-Jul	60 95	61 60	130 210	20 30	98* 158*
American River Folsom Reservoir	Apr-Jul	710	58	1600	230	1230


^{*30} Year Averages for 1971-2000 are incomplete. Those forecast points with an asterisk have incomplete averages, so 1961-1990 averages are listed. The new averages will be incorporated into this report when the complete data sets become available.

^{**} Pit River 30-year average is full natural flow.


Sacramento/Trinity/Klamath River Basins

Seasonal Basin Precipitation

October 1 to Date




Seasonal Basin Snowpack Water Content in % of April 1 Average




Sacramento/Trinity/Klamath River Basins


Basin Reservoir Storage

Contents of Major Reservoirs in % of Average



Seasonal Basin Runoff October 1 to Date






SAN JOAQUIN BASIN

		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
South Fork San Joaquin River Hooper Ck, blo, Florence Lk, nr	Apr-Jul	115	60	240	60	192*
San Joaquin River Millerton Lake	Apr-Jul	760	60	1550	300	1270
Merced River Pohono Bridge, at, Yosemite, nr Merced Falls, blo	Apr-Jul Apr-Jul	235 355	65 55	450 760	90 130	360* 645
Tuolumne River Hetch Hetchy, nr La Grange, nr	Apr-Jul Apr-Jul	385 740	65 60	740 1500	160 310	596* 1230
Middle Fork Stanislaus River Beardsley Dam, blo	Apr-Jul	190	59	400	80	320*
Stanislaus River New Melones Dam	Apr-Jul	380	55	830	140	695
North Fork Mokelumne River West Point	Apr-Jul	230	55	500	80	416*
Mokelumne River Pardee Reservoir	Apr-Jul	250	54	535	105	460
Cosumnes River Michigan Bar	Apr-Jul	65	53	180	10.0	123


^{*30} Year Averages for 1971-2000 are incomplete. Those forecast points with an asterisk have incomplete averages, so 1961-1990 averages are listed. The new averages will be incorporated into this report when the complete data sets become available.

San Joaquin Basin

Seasonal Basin Precipitation October 1 to Date




Seasonal Basin Snowpack Water Content in % of April 1 Average




San Joaquin Basin

Basin Reservoir Storage Contents of Major Reservoirs in % of Average



Season Basin Runoff October 1 to Date




TULARE LAKE BASIN

		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
Kern River						
Kernville, nr	Apr-Jul	260	65	480	100	398*
Isabella Dam, blo	Apr-Jul	290	60	580	110	480
Bakersfield, nr	Apr-Jul	295	60	600	115	490
Tule River						
Success Dam	Apr-Jul	30	45	80	10.0	66
Kaweah River						
Terminus Dam	Apr-Jul	175	60	350	70	290
North Fork Kings River						
Cliff Camp, nr	Apr-Jul	145	60	290	60	240*
Kings River						
Pine Flat Dam, blo	Apr-Jul	750	60	1500	300	1250


^{*30} Year Averages for 1971-2000 are incomplete. Those forecast points with an asterisk have incomplete averages, so 1961-1990 averages are listed. The new averages will be incorporated into this report when the complete data sets become available.

Tulare Lake Basin

Seasonal Precipitation October 1 to Date




Seasonal Basin Snowpack Water Content in % of April 1 Average





Tulare Lake Basin

Basin Reservoir Storage Contents of Major Reservoirs in % of Average



Seasonal Basin Runoff October 1 to Date






EAST SIDE SIERRA NEVADA BASINS


		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
Truckee River						
Truckee River Lake Tahoe Stage Rise	Apr-High	0.70	51	1.84	0.04	1.38
Little Truckee River Stampede Dam	Apr-Jul	39	49	121	2.4	80
Truckee River Farad	Apr-Jul	125	48	390	7.8	260
Carson River						
East Fork Carson River Gardnerville, nr	Apr-Jul	95	50	215	5.7	189
West Fork Carson River Woodfords	Apr-Jul	28	50	69	2.2	56
Carson River Carson City, nr Fort Churchill, nr	Apr-Jul Apr-Jul	70 55	37 31	255 191	1.88	188 178
Walker River						
East Walker River Bridgeport, nr	Apr-Aug	32	48	84	2.0	67
West Walker River Ltl Walker, blo, Coleville, nr	Apr-Jul	85	54	189	6.2	156

East Side Sierra Nevada Basins

Seasonal Basin Precipitation October 1 to Date




Basin Snowpack % of Average SWE to Date




East Side Sierra Nevada Basins

Seasonal Basin Runoff

October 1 to Date



Humboldt River Basin




Water Supply Forecasts

		Most Prob Vol KAF	Most Prob Vol %Norm	Reas Max Vol KAF	Reas Min Vol KAF	30 Year Avg KAF
North Fork Humboldt River Devils Gate, at, Halleck, nr	Apr-Jul	14.0	41	42	0.68	34*
South Fork Humboldt River Dixie Ck, abv, Elko, nr	Apr-Jul	36	47	107	3.8	76
Marys River Hot Springs, abv, Deeth, nr	Apr-Jul	15.0	38	38	1.17	39
Humboldt River Elko, nr Palisade Comus Imlay, nr	Apr-Jul Apr-Jul Apr-Jul Apr-Jul	55 90 60 40	36 36 27 21	146 195 184 185		154 250 225 188
Martin Ck Paradise Vly, nr	Apr-Jul	9.0	0 48	26	0.37	18.7

^{*30} Year Averages for 1971-2000 are incomplete. Those forecast points with an asterisk have incomplete averages, so 1961-1990 averages are listed. The new averages will be incorporated into this report when the complete data sets become available.

Humboldt River Basin

Seasonal Basin Precipitation October 1 to Date



Basin Snowpack % of Average SWE to Date

